

## Chapter 55


## ACID-BASE EQUILIBRIA, Part 2

©2012, 2023 Mark E. Noble

We now turn to calculations involving weak acid equilibria. We've seen such a calculation already: Example 4 in Chapter 52 involved the dissociation of acetic acid. That's very typical of the calculations which you will be doing here. We will now add pH and we will step up our approximation methods. We will also consider acids with more than one dissociation step. Despite these extras, the fundamentals remain the same as done previously. After the calculations, we'll talk a bit about why some acids are strong or weak.

55.1  $K_a$ 

Consider a generic weak acid, HA, undergoing dissociation.




The equilibrium constant  $K$  for acid dissociation is specifically designated  $K_a$ .

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

Values of  $K_a$  have been measured for many acids and these vary widely. Common values are in the range of  $10^{-2}$  to  $10^{-10}$  although some are outside that. As the value of  $K_a$  decreases, there is less dissociation and less  $\text{H}^+$  at equilibrium, and this constitutes a weaker acid. Conversely, a better acid will be reflected by a larger  $K_a$ . A selection of  $K_a$  values is given in Appendix B.

A weak acid produces some  $\text{H}^+$  in water and this amount of  $\text{H}^+$  is in addition to some  $\text{H}^+$  which is produced by autoionization. For a typical problem involving a weak acid, we are concerned with the  $[\text{H}^+]$  and the pH of some solution, and that could require consideration of both of these sources for total  $[\text{H}^+]$ . Fortunately, autoionization has only a very small impact on total  $[\text{H}^+]$  in most solutions of acids and bases, and we can therefore ignore it. The reason for this lies in equilibrium dynamics. The autoionization equilibrium



has a very small  $K_w = 1.0 \times 10^{-14}$ , from which  $[\text{H}^+] = [\text{OH}^-] = 1.0 \times 10^{-7}$  M in pure water. Upon adding weak acid, the  $\text{H}^+$  from the acid shifts the above equilibrium to the left, which suppresses the autoionization. For a quantitative illustration, consider some weak acid at  $[\text{HA}] = 0.010$  M and with  $K_a = 1.0 \times 10^{-6}$ . In this case, the weak acid dissociation by itself provides  $[\text{H}^+] = 1.0 \times 10^{-4}$  M. If we plug this value for  $[\text{H}^+]$  into  $K_w$

$$K_w = [\text{H}^+][\text{OH}^-] = (1.0 \times 10^{-4})[\text{OH}^-] = 1.0 \times 10^{-14}$$

then we find  $[\text{OH}^-] = 1.0 \times 10^{-10}$  M. The autoionization process by itself gives equal amounts of  $\text{H}^+$  and  $\text{OH}^-$ , which means autoionization's contribution to total  $[\text{H}^+]$  is likewise  $1.0 \times 10^{-10}$  M. These numbers are a thousand-fold less compared to the concentrations in pure water, and this shows the shift to the left for autoionization. The grand total for  $[\text{H}^+]$  is now the sum of the amounts from acid dissociation and from autoionization.

$[\text{H}^+]$ from weak acid dissociation:	0.00010 M
$[\text{H}^+]$ from autoionization:	<u>0.00000000010 M</u>
Grand total $[\text{H}^+]$ :	0.00010 M

The result is that  $[\text{H}^+]$  from autoionization is not significant to total  $[\text{H}^+]$  when the acid is present. For our coverage, this will be general and we will ignore the autoionization part completely. Although there are cases close to neutral where autoionization can still be significant, we will not deal with those. This discussion also applies to  $\text{OH}^-$  in the presence of added bases. Thus, when we get to bases, we will ignore autoionization at that time also.

OK, let's start.

.....

**Example 1.** The term hydrohalic acid applies to a solution of any of the hydrogen halides dissolved in water. Unlike HCl, HBr and HI which are strong, HF is weak with  $K_a = 6.8 \times 10^{-4}$ . A solution of HF is prepared by dissolving 0.0448 mol HF in water to make 200. mL of solution. What are the concentrations at equilibrium of HF,  $\text{H}^+$  and  $\text{F}^-$ ? What are the pH of the solution and the percent dissociation of HF?

.....

Except for pH, this is the same type of problem as Example 4 in Chapter 52. Look over that again if you need to get back in the groove.



► Step 2.  $K_a$

$$K_a = \frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]} = 6.8 \times 10^{-4}$$

► Step 3. We need an initial amount and a table to put it in.

$$\text{initial } [\text{HF}] = \frac{0.0448 \text{ mol}}{0.200 \text{ L}} = 0.224 \text{ M}$$

	[HF]	[H <sup>+</sup> ]	[F <sup>-</sup> ]
Initial:	0.224	-0-	-0-

► Step 4. Bring in the changes.

	[HF]	[H <sup>+</sup> ]	[F <sup>-</sup> ]
Initial:	0.224	-0-	-0-
Changes:	-x	+x	+x

► Step 5. Final equilibrium amounts.

	[HF]	[H <sup>+</sup> ]	[F <sup>-</sup> ]
Initial:	0.224	-0-	-0-
Changes:	-x	+x	+x
Equilibrium:	0.224 - x	x	x

► Step 6. Plug into  $K_a$ .

$$K_a = \frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]} = \frac{(x)(x)}{0.224 - x} = 6.8 \times 10^{-4}$$

Re-arrange.

$$x^2 + 6.8 \times 10^{-4}x - 1.5 \times 10^{-4} = 0$$

Solve for  $x$  using the quadratic equation. You will get  $x = -0.013$  and  $0.012$ . The negative is the nonsense answer.

► Step 7. Take 0.012 back into the equilibrium line of the table for the final concentrations.

$$\begin{aligned} [\text{HF}] &= (0.224 - 0.012) \text{ M} = 0.212 \text{ M} \\ [\text{H}^+] &= 0.012 \text{ M} \\ [\text{F}^-] &= 0.012 \text{ M} \end{aligned}$$

► Step 8.  $K$ -check gives  $6.8 \times 10^{-4}$ , right on.

Time out. We threw this into quadratic but what about approximation? Would this have worked?  $K_a$  is getting close to  $10^{-3}$  which is not real small, so this could be iffy. Let's go ahead and try it. Assume we haven't done the quadratic above. Start over at  $K_a$ .

$$K_a = \frac{(x)(x)}{0.224 - x} = 6.8 \times 10^{-4}$$

If we assume  $x$  is small relative to 0.224, then we approximate  $0.224 - x \approx 0.224$ . That gives us

$$K_a \approx \frac{(x)(x)}{0.224} = 6.8 \times 10^{-4}$$

which leads to  $x = \pm 0.012$ . Only the positive result is valid and it agrees with the value from the quadratic equation, so approximation works here also.

OK, we've got  $x$  and we've got the concentrations. We still need a pH and a percent dissociation.


$$\text{pH} = -\log(0.012) = 1.92$$

$$\% \text{diss} = \frac{\text{amount which dissociated}}{\text{starting amount}} \times 100\% = \frac{0.012 \text{ M}}{0.224 \text{ M}} \times 100\% = 5.4\%$$

We're done.

Approximation worked fine in this Example, but I will tell you that it's close to not working. Now we will take our approximation methods up one notch and bring in another consideration. In Chapter 52, we considered the size of  $K$  as an indicator of whether approximation might work or not. We will now add concentrations as another indicator. Recall from the Second Principle of Equilibrium Dynamics that dilution favors more  $Q$  components. In Chapter 53, we considered this for  $\text{CH}_3\text{CO}_2\text{H}$  dissociation.

“ Apply this to the acetic acid equilibrium.




If we add water, the immediate effect is to decrease all concentrations and decrease  $Q$ . In order to return to equilibrium,  $[\text{CH}_3\text{CO}_2^-]$  and  $[\text{H}^+]$  will increase while  $[\text{CH}_3\text{CO}_2\text{H}]$  will decrease. In the end, more moles of  $\text{CH}_3\text{CO}_2\text{H}$  are actually dissociated at the new point of balance. The net result is that the percent dissociation increases upon dilution. This is an extremely important notion in many solution equilibria, as will be seen later in Chapter 55. ”

Later is now. For all dissociation equilibria, dilution favors more dissociation. Know that. But what's that got to do with approximation? More dissociation means a bigger relative shift and that can spell trouble for approximation. We will illustrate this dilution dilemma by taking Example 1 and diluting it 20-fold.

**Example 2.** Repeat Example 1 but with a total solution volume of 4.00 L. What are the concentrations at equilibrium of HF,  $\text{H}^+$  and  $\text{F}^-$ ? What are the pH of the solution and the percent dissociation of HF?

The setup is the same but some numbers are different. The balanced equation and  $K_a$  are the same.



$$K_a = \frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]} = 6.8 \times 10^{-4}$$

You need a new initial

$$\text{initial } [\text{HF}] = \frac{0.0448 \text{ mol}}{4.00 \text{ L}} = 0.0112 \text{ M}$$

and a couple of changes to the table.

	[HF]	[H <sup>+</sup> ]	[F <sup>-</sup> ]
Initial:	0.0112	-0-	-0-
Changes:	-x	+x	+x
Equilibrium:	0.0112 - x	x	x

Plug into  $K_a$ .

$$K_a = \frac{[\text{H}^+][\text{F}^-]}{[\text{HF}]} = \frac{(x)(x)}{0.0112 - x} = 6.8 \times 10^{-4}$$

Re-arrange.

$$x^2 + 6.8 \times 10^{-4} x - 7.6 \times 10^{-6} = 0$$

The quadratic equation will give  $x = -0.0031$  and  $0.0024$ . The negative is again nonsense. Put  $0.0024$  into the last line of the table for  $x$ .

$$\begin{aligned} [\text{HF}] &= (0.0112 - 0.0024) \text{ M} = 0.0088 \text{ M} \\ [\text{H}^+] &= 0.0024 \text{ M} \\ [\text{F}^-] &= 0.0024 \text{ M} \end{aligned}$$

You can do the  $K$ -check. It's close enough.

Now we pause to bring in approximation for comparison. Again, assume we don't know the answers from the quadratic solution. Let's again go back to  $K_a$ .

$$K_a = \frac{(x)(x)}{0.0112 - x} = 6.8 \times 10^{-4}$$

Assume  $x$  is small relative to 0.0112; thus,  $0.0112 - x \approx 0.0112$ .

$$K_a \approx \frac{(x)(x)}{0.0112} = 6.8 \times 10^{-4}$$

Solving for  $x$  will now give  $\pm 0.0028$  and the negative is again nonsense. Unfortunately, the positive 0.0028 is now a bit different from the answer by the quadratic route, so the present answer is not good enough. Approximation does not work well here. Too bad.

Let's finish off this Example using 0.0024 from the quadratic result. The pH is

$$\text{pH} = -\log(0.0024) = 2.62$$

and the percent dissociation is

$$\% \text{diss} = \frac{\text{amount which dissociated}}{\text{starting amount}} \times 100\% = \frac{0.0024 \text{ M}}{0.0112 \text{ M}} \times 100\% = 21\%$$

and that wraps up the final answers.

Note the 21% dissociation. That's what killed simple approximation here. That 21% is way too much of a relative change. In Example 1, the relative change was only 5.4% and approximation worked. The  $K_a$  value was the same for both Examples, so this illustrates that a  $K$  by itself is not the only indicator to consider. We now refine our approach as follows. For dissociation equilibria, approximation works best for smaller  $K$  and/or higher concentrations. Both of these factors give less relative change (smaller %diss) from initial conditions. Note also the converse. When  $K$  is moderate (not so small) and/or when concentrations are more dilute, then approximation is more prone to fail.

So does that mean the 0.0028 answer from simple approximation is totally wrong? That's up to your instructor. That's not the end of approximation, however, because there is a refinement available called iteration.

### 55.2 Iterate, iterate, iterate.

The term "iterate" follows the normal dictionary meaning which is to repeat. Here, iteration involves repeat approximation. There are four steps to iteration but the first is not new.

#### Iteration

- Do the simple approximation for  $x$ .
- Plug this answer into the quantity which was approximated.
- Solve for a new  $x$ .
- Repeat Steps B and C until consecutive answers agree to the correct sigfigs (or cycle within the last sigfig, although this is less common).

For Step B, the underlined phrase is important. For Step D, the parenthetical part will be illustrated in Example 3.

Let's illustrate iteration using Example 2. Upstairs, we did the usual, simple approximation

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112} = 6.8 \times 10^{-4}$$

which gave  $x = 0.0028$ . That ends Step A of iteration. For Step B, we plug this value for  $x$  back into the quantity which we had approximated. We only approximated the denominator; we did not approximate the numerator. Thus, we plug in 0.0028 for  $x$  in the denominator only.

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112 - 0.0028} = \frac{x^2}{0.0084} = 6.8 \times 10^{-4}$$

For Step C, solve again for  $x$ : you get 0.0024. That's quite different from 0.0028. Are we done? Step D says do Steps B and C again.

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112 - 0.0024} = \frac{x^2}{0.0088} = 6.8 \times 10^{-4}$$

Solving gives  $x = 0.0024$  which is the same as the prior result. Now Step D says stop. This is your final answer for  $x$ . Notice that it matches the value from the full quadratic solution done previously. Iteration worked.

So what? The full quadratic worked also. Why do we need iteration?

Good question. It's another tool, and it's a very good tool in a range of applications. Iteration methodologies can get very sophisticated; they play a critically important role in many computational methods, in which case the repetitive part is run by software. Those programs can run iterations through hundreds or thousands or more cycles. While our use here is not that sophisticated, you can still get a feel for the method. For our types of problems, iteration does three things. First, it provides a check on the usual approximation. Thus, if you are in doubt whether simple approximation is good enough for a particular problem, then iteration will tell you. Second, iteration gives more leeway to the range of  $K$ 's and concentrations which can be done, as demonstrated by its success in handling Example 2 when simple approximation was not enough. Third, not all problems can be solved directly and some form of approximation will be necessary in those cases.

Although it is better than simple approximation and it gives more leeway, iteration is still an approximation method and it should not be used where it doesn't belong. A notable feature for iteration is that the differences between successive values for  $x$  should be getting smaller, eventually going to zero. If successive iterations for  $x$  are jumping around with no clear pattern, then iteration may be failing. Also, iteration is intended to simplify the math in many cases. If you're doing 10 or 12 or more iterations, then a direct algebraic solution (such as by way of the quadratic equation) may have been much faster. Of course, there's always the possibility of a math error somewhere, which can happen with any method. Be ready to check your math.

In our current usage for a weak acid dissociation problem, you now have two tools to choose from. You can solve directly by quadratic equation or you can solve by approximation/iteration. The two methods usually give the same answer although sometimes they can differ by one in the last sigfig. That's no big deal and either answer would be acceptable. Besides, we are sort of taking it easy on the quadratic equation anyway with our simplified sigfig rule. So, use either way; it's your call. Of course, if your instructor wants them done a certain way, then call it that way. But if you have the choice, then you need to decide which way is faster and more reliable for you. Try them both and find out. Here: compare the two methods for Example 2 starting from the  $K_a$  expression.

$$K_a = \frac{(x)(x)}{0.0112 - x} = 6.8 \times 10^{-4}$$

For direct (quadratic) method, you had to get from there to

$$x^2 + 6.8 \times 10^{-4} x - 7.6 \times 10^{-6} = 0$$

and then into

$$x = \frac{-(-6.8 \times 10^{-4}) \pm \sqrt{(6.8 \times 10^{-4})^2 - 4(1)(-7.6 \times 10^{-6})}}{2(1)}$$

in order to get  $x$ . For the iteration method, you worked with the  $K_a$  expression directly and you had to execute three cycles.

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112} = 6.8 \times 10^{-4} \quad \rightarrow \rightarrow \rightarrow x = 0.0028$$

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112 - 0.0028} = \frac{x^2}{0.0084} = 6.8 \times 10^{-4} \quad \rightarrow \rightarrow \rightarrow x = 0.0024$$

$$K_a = \frac{x^2}{0.0112 - x} \approx \frac{x^2}{0.0112 - 0.0024} = \frac{x^2}{0.0088} = 6.8 \times 10^{-4} \quad \rightarrow \rightarrow \rightarrow x = 0.0024$$

Three cycles are not a lot, but some problems will run more. It just depends. So, overall, which method would you prefer? Some students can rip through an algebraic re-arrangement and quadratic solution in no time. Other students are more prone to errors in the middle steps of the derivation and tend to shy away from it. On the other hand, iteration can be tedious, especially if a lot of cycles are involved. You

have to try both ways. Another plus for iteration is that it can help in other types of problems in chemistry and also in other fields.

Moving on, here's another Example and a quirk. The quirk relates to the parenthetical part of Iteration Step D above.

**Example 3.** 0.00820 mol  $\text{HNO}_2$  was dissolved in a solution of total volume 200.0 mL. At equilibrium, what are  $[\text{H}^+]$  and the pH?  $K_a$  for nitrous acid is  $7.1 \times 10^{-4}$ .

Balanced equation  $\text{HNO}_2(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{NO}_2^-(\text{aq})$

$K_a$  expression

$$K_a = \frac{[\text{H}^+][\text{NO}_2^-]}{[\text{HNO}_2]} = 7.1 \times 10^{-4}$$

Set up the initial amount.

$$\text{initial } [\text{HNO}_2] = \frac{0.00820 \text{ mol}}{0.200 \text{ L}} = 0.0410 \text{ M}$$

Now set the table.

	$[\text{HNO}_2]$	$[\text{H}^+]$	$[\text{NO}_2^-]$
Initial:	0.0410	-0-	-0-

Add changes.

	$[\text{HNO}_2]$	$[\text{H}^+]$	$[\text{NO}_2^-]$
Initial:	0.0410	-0-	-0-
Changes:	-x	+x	+x

Add final equilibrium amounts.

	$[\text{HNO}_2]$	$[\text{H}^+]$	$[\text{NO}_2^-]$
Initial:	0.0410	-0-	-0-
Changes:	-x	+x	+x
Equilibrium:	$0.0410 - x$	x	x

Plug into  $K_a$ .

$$K_a = \frac{[\text{H}^+][\text{NO}_2^-]}{[\text{HNO}_2]} = \frac{(x)(x)}{0.0410 - x} = 7.1 \times 10^{-4}$$

Solve for x. We'll first do the quadratic route. Re-arrange.

$$x^2 + 7.1 \times 10^{-4} x - 2.9 \times 10^{-5} = 0$$

You will get  $x = 0.0050$  as the usable answer.

Now solve for x using iteration. For this, approximate the final equilibrium value for  $[\text{HNO}_2]$  as  $0.0410 - x \approx 0.0410$ .

$$K_a = \frac{x^2}{0.0410 - x} \approx \frac{x^2}{0.0410} = 7.1 \times 10^{-4}$$

Solving for x gives  $x = 0.0054$ . Iterate.

$$K_a = \frac{x^2}{0.0410 - x} \approx \frac{x^2}{0.0410 - 0.0054} = \frac{x^2}{0.0356} = 7.1 \times 10^{-4}$$

This gives  $x = 0.0050$ . Iterate again.

$$K_a = \frac{x^2}{0.0410 - x} \approx \frac{x^2}{0.0410 - 0.0050} = \frac{x^2}{0.0360} = 7.1 \times 10^{-4}$$

$x = 0.0051$ . Go again.

$$K_a = \frac{x^2}{0.0410 - x} \approx \frac{x^2}{0.0410 - 0.0051} = \frac{x^2}{0.0359} = 7.1 \times 10^{-4}$$

You again get  $x = 0.0050$ , which was the same value as two  $x$ 's ago. If you iterate that, you will get 0.0051, and these two numbers just cycle back and forth with each iteration. This is somewhat of a quirk and it ties into how we are dealing with the sigfigs. This is not a very common outcome. If it does happen to you on some problem, just stop when you repeat a prior number and average the two repeating numbers together. Here, averaging 0.0050 and 0.0051 gives 0.0050 to the correct sigfigs. This does match the value from the quadratic route but a difference of 0.0001 would have been OK anyway.

The  $K$  checks out fine either way. In conclusion, we have  $x = 0.0050$  as the molarity for  $[H^+]$ . This gives  $\text{pH} = 2.30$ .

Your turn. By the way, from now on, if the  $K_a$  is not given in the problem, check Appendix B.

.....  
**Example 4.** 0.0181 mol of hypochlorous acid is dissolved in 500. mL solution. What are the concentrations at equilibrium of HClO,  $H^+$  and  $ClO^-$ ? What are the pH and the percent dissociation?  
 .....

Balanced equation:

$K_a$  expression:

Initial amount:

Fill in a table:

	[HClO]	[H <sup>+</sup> ]	[ClO <sup>-</sup> ]
Initial:			
Changes:			
Equilibrium:			

Plug into  $K_a$ .

How do you want to solve for  $x$ ? Take your pick. Better yet, do them both.

Quadratic:

Approximation/iteration:

Once you've got  $x$ , you can do the concentrations.

[HClO] =

[H<sup>+</sup>] =

$$[\text{ClO}^-] =$$

You can do your  $K$ -check.

If that's OK, do the pH,

and, finally, do the percent dissociation.

As one check for your answers, the pH is 4.48. The very small percent dissociation made it very straightforward for approximation/iteration.


These Examples 1 - 4 show the typical calculations and methods which are involved in many acid dissociation problems. Of course, variations are possible. Here's one.

.....

**Example 5.** Phenol,  $\text{C}_6\text{H}_5\text{OH}$ , was mentioned back in Section 36.1 for sublimation, smell, and antiseptic applications. In water it is a weak acid, and historically it was known as carbolic acid. 0.0136 mol of phenol is dissolved in 100. mL solution. The pH at equilibrium is measured to be 5.43. Find  $K_a$  and  $\Delta G^\circ$  for the dissociation of phenol at 25 °C.

.....

This starts out very differently and it asks for different things. Go ahead and set up a balanced equation and a  $K_a$  expression.



and then initialize your acid.

$$\text{initial } [\text{C}_6\text{H}_5\text{OH}] = 0.136 \text{ M}$$

Now what?

For this Example, you must find the value for  $K_a$ . (It's not in Appendix B.) That will then lead to  $\Delta G^\circ$ . We could find  $K_a$  if we have some numbers to plug in for equilibrium concentrations, but we have an initial (not equilibrium) acid concentration. Note that the pH was given at equilibrium, so we can get the equilibrium amount of  $[\text{H}^+]$  from the pH and the other concentrations from that. Since these are at equilibrium, this is not a change problem and we don't need a change table.

$$[\text{H}^+] = 10^{-5.43} \text{ M} = 3.7 \times 10^{-6} \text{ M}$$

The dissociation gives  $\text{H}^+$  and  $\text{C}_6\text{H}_5\text{O}^-$  in equal amounts.

$$[\text{H}^+] = 3.7 \times 10^{-6} \text{ M} = [\text{C}_6\text{H}_5\text{O}^-]$$

The equilibrium amount of  $[\text{C}_6\text{H}_5\text{OH}]$  is the initial amount minus the amount which has dissociated.

$$[\text{C}_6\text{H}_5\text{OH}] = 0.136 \text{ M} - 3.7 \times 10^{-6} \text{ M} = 0.136 \text{ M}$$

Now you've got everything for  $K_a$ .

$$K_a = \frac{[\text{H}^+][\text{C}_6\text{H}_5\text{O}^-]}{[\text{C}_6\text{H}_5\text{OH}]} = \frac{(3.7 \times 10^{-6})(3.7 \times 10^{-6})}{0.136} = 1.0 \times 10^{-10}$$

Use that to find  $\Delta G^\circ$ .

$$\Delta G^\circ = -RT \ln K_a = -(8.314 \text{ J/K})(298 \text{ K}) \ln(1.0 \times 10^{-10}) = 57 \text{ kJ}$$

The value is positive and the process is endergonic, as to be expected for a weak acid.

### 55.3 Mono vs. poly


Let's now dissociate a bit more.

All calculations with weak acids so far have involved monoprotic acids. A monoprotic acid is an acid which is capable of losing one  $\text{H}^+$ . Acids which can lose more than one  $\text{H}^+$  are called polyprotic acids.




Within that category, these can be further specified by the number of  $H^+$  which can be lost. For example, a diprotic acid can lose two  $H^+$  and a triprotic can lose three.  $H_2SO_4$  is the most common example of a diprotic while  $H_3PO_4$  is the most common example of a triprotic. Other examples of diprotics are  $H_2CO_3$  and  $H_2C_2O_4$ , while another triprotic is given by citric acid,  $C_3H_5O(CO_2H)_3$ , of citrus fruit fame. Some acids can lose four or more  $H^+$  and the terms do go higher than triprotic, but those are much less common.

Polyprotic acids often undergo dissociation one step at a time and each step is an equilibrium. Each step takes a number, and the first dissociation begins with the starting, neutral acid. Here is an illustration of the two steps for oxalic acid.




$$K_{a1} = \frac{[H^+][HC_2O_4^-]}{[H_2C_2O_4]} = 0.054$$




$$K_{a2} = \frac{[H^+][C_2O_4^{2-}]}{[HC_2O_4^-]} = 5.4 \times 10^{-5}$$

Note the inclusion of subscript numbers within the  $K_a$ 's to indicate the step number. You need to be careful with step numbers. Whenever a number for a dissociation is given, then it traces back to starting, neutral acid; the dissociation of the starting, neutral acid is the first step. You don't always have to work with step numbers, however, as long as you are careful with the meaning. For example, we can independently consider the dissociation of  $HC_2O_4^-$  by itself




and the  $K_a$  for this equation is the  $K_a$  for  $HC_2O_4^-$ . This equation is the same as the second dissociation of  $H_2C_2O_4$ . Thus, " $K_a$  for  $HC_2O_4^-$ " is the same as saying " $K_{a2}$  for  $H_2C_2O_4$ ". Watch the wording.

Carbonic acid,  $H_2CO_3$ , is also diprotic but this one is not straightforward.  $H_2CO_3$  has only a feeble existence in aqueous solution. This was first noted back in Section 11.4, and we can now elaborate more upon the equilibria involved. When  $CO_2(g)$  dissolves in water, some  $H_2CO_3(aq)$  is produced but the reality is that very little  $H_2CO_3$  is present at equilibrium; most of the solute is simply  $CO_2(aq)$ . Carbonic acid can be made as a pure compound but, in water, it decomposes to  $CO_2(aq)$  and  $H_2O(l)$ .



$$K(\text{decomp}) = \frac{[CO_2]}{[H_2CO_3]}$$

It has been difficult to measure  $K(\text{decomp})$  precisely, and values in the range of 380 - 800 have been reported. Nevertheless, the bottom line is that dissolved  $CO_2$  is mostly  $CO_2(aq)$ , and less than 0.3% is  $H_2CO_3$ . The primary equilibria for acid dissociation are therefore based on  $CO_2(aq)$ .



$$K_{a1} = \frac{[H^+][HCO_3^-]}{[CO_2]} = 4.5 \times 10^{-7}$$



$$K_{a2} = \frac{[H^+][CO_3^{2-}]}{[HCO_3^-]} = 4.7 \times 10^{-11}$$

Although the amount of  $CO_2$  greatly exceeds the amount of  $H_2CO_3$ , you can still consider the dissociation of  $H_2CO_3$  directly. This will likewise be two steps. The first step looks typical and the second step is the same as above.




$$K_{a1} = \frac{[H^+][HCO_3^-]}{[H_2CO_3]}$$



$$K_{a2} = \frac{[H^+][CO_3^{2-}]}{[HCO_3^-]} = 4.7 \times 10^{-11}$$

The value of  $K_{a1}$  for  $\text{H}_2\text{CO}_3$  is also difficult to measure. It is actually related to  $K(\text{decomp})$  by summation of the following equilibria.




Since we are summing equations, we multiply  $K$ 's.

$$K(\text{decomp}) \times K_{a1}(\text{CO}_2) = K_{a1}(\text{H}_2\text{CO}_3)$$

$K_{a1}$  for  $\text{CO}_2$  is known very well ( $4.5 \times 10^{-7}$ ), but the others are not. From the range of 380 - 800 cited above for  $K(\text{decomp})$ , you can calculate  $K_{a1}(\text{H}_2\text{CO}_3)$  to be in the range of  $1.7 \times 10^{-4}$  -  $3.6 \times 10^{-4}$ . Due to the difficulty of measuring  $K(\text{decomp})$  and  $K_{a1}(\text{H}_2\text{CO}_3)$ , calculations for these solutions are simply based on  $\text{CO}_2$  and its  $K_{a1}$ .

$\text{SO}_2$  was also noted back in Section 11.4 for similarities to  $\text{CO}_2$  but there are also some differences.  $\text{SO}_2$  was thought for many years to dissolve in water to produce sulfurous acid,  $\text{H}_2\text{SO}_3$ , but that appears to be wrong. Unlike  $\text{H}_2\text{CO}_3$  which does have some minor existence in water, there is no  $\text{H}_2\text{SO}_3$  to any measurable extent. In fact,  $\text{H}_2\text{SO}_3$  has never even been prepared in pure form under any condition. Thus, " $\text{H}_2\text{SO}_3(aq)$ " really means  $\text{SO}_2(aq)$  and  $\text{H}_2\text{O}(l)$ . The weak acid steps are the following.




$$K_{a1} = \frac{[\text{H}^+][\text{HSO}_3^-]}{[\text{SO}_2]} = 0.014$$



$$K_{a2} = \frac{[\text{H}^+][\text{SO}_3^{2-}]}{[\text{HSO}_3^-]} = 6.7 \times 10^{-8}$$

Since there is no  $\text{H}_2\text{SO}_3$  in water, then these are the only equilibria to consider.

The cases of  $\text{CO}_2(aq)$  and  $\text{SO}_2(aq)$  are different from other acids but that alone would not warrant their mention here. What does warrant their mention is their tremendous importance on a huge scale. For example,  $\text{CO}_2$  and  $\text{SO}_2$  are immensely important in the chemistry of Earth's atmosphere, as gases and as solutes in air-borne water droplets. Both contribute in a big way to acid rain. Although there is far more  $\text{CO}_2$  in the atmosphere,  $\text{SO}_2$  has a much greater  $K_{a1}$ . Furthermore,  $\text{SO}_2$  is oxidized in the atmosphere and eventually converted to the strong acid,  $\text{H}_2\text{SO}_4$ . Besides pH effects,  $\text{CO}_2$  is also a major greenhouse gas which adds to its importance in global warming. But the impact of  $\text{CO}_2$  is not just up in the air.  $\text{CO}_2$  is of huge physiological importance to humans, to other animals, and to plants. The interplay of  $\text{CO}_2(g)$ ,  $\text{CO}_2(aq)$ ,  $\text{H}_2\text{CO}_3(aq)$ ,  $\text{HCO}_3^-(aq)$  and  $\text{CO}_3^{2-}(aq)$  at physiological pH are all vastly important to various biological systems. The aqueous equilibria above play an important role in these processes, as does the solubility equilibrium for  $\text{CO}_2(g)$  which we have considered at various places in prior Chapters.




There's no escaping it: these things are part of your world. A **BIG** part.

By the way, as a practical matter, the peculiarities of  $\text{CO}_2$  and  $\text{SO}_2$  do not change a typical  $K_a$  or pH calculation. You just treat the acidity of  $\text{CO}_2$  and of  $\text{SO}_2$  as you would any other weak acid, given some initial amounts for  $\text{CO}_2$  or  $\text{SO}_2$ . We'll cover a  $\text{CO}_2$  problem in Example 6 below but let me finish off a few other points first.

The polyprotic dissociations so far are for diprotic acids. Let's now take a look at a triprotic,  $\text{H}_3\text{PO}_4$ .



$$K_{a1} = \frac{[\text{H}^+][\text{H}_2\text{PO}_4^-]}{[\text{H}_3\text{PO}_4]} = 7.1 \times 10^{-3}$$



$$K_{a2} = \frac{[\text{H}^+][\text{HPO}_4^{2-}]}{[\text{H}_2\text{PO}_4^-]} = 6.3 \times 10^{-8}$$




$$K_{a3} = \frac{[\text{H}^+][\text{PO}_4^{3-}]}{[\text{HPO}_4^{2-}]} = 4.5 \times 10^{-13}$$

As always, for each step of dissociation, you lose one  $\text{H}^+$ . Also, keep in mind the part about step numbers. For example, note that  $K_a$  for  $\text{HPO}_4^{2-}$  is the same as  $K_{a3}$  of  $\text{H}_3\text{PO}_4$ .

In the polyprotic series cited so far,  $\text{HC}_2\text{O}_4^-$ ,  $\text{HCO}_3^-$ ,  $\text{HSO}_3^-$ ,  $\text{H}_2\text{PO}_4^-$  and  $\text{HPO}_4^{2-}$  are all hydrogen anions. All hydrogen anions derive from polyprotic acids and all are middle members of the polyprotic series. So far, all of these hydrogen anions are amphoteric.

Let's now consider  $\text{H}_2\text{SO}_4$ . This is different from most polyprotics because its first dissociation is strong. Although the first step is strong, its second dissociation is weak.



$$K_{a2} = \frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} = 0.010$$

$K_{a1}$  is not shown because  $K_a$ 's are usually not well known for strong acids. The first dissociation is strong because  $\text{HSO}_4^-$  is not pulling back on  $\text{H}^+$  to a significant extent. Since it is not pulling back,  $\text{HSO}_4^-$  is not basic. Since it is not basic,  $\text{HSO}_4^-$  is not amphoteric; it is only acidic. This will be general:

All hydrogen anions derived from strong acids are acidic but not amphoteric.

This connects to your flag in Section 54.4. Remember overall that hydrogen anions can be amphoteric or (only) acidic.

Let's start in on some calculations.

Multiple dissociation steps can cause difficulty in calculations involving polyprotics. Since each step can produce some  $\text{H}^+$ , the question arises as to whether we need to worry about all of the dissociation steps for total  $[\text{H}^+]$ . The answer is: it depends.



For polyprotics whose first step is strong, then the calculation for total  $\text{H}^+$  must involve both the strong first step and the weak second step. We will see this for  $\text{H}_2\text{SO}_4$  in Example 7. For polyprotic acids with all weak steps, then the calculation for total  $\text{H}^+$  can get ugly. To avoid utter ugliness, we impose a simplification: we limit coverage to calculations where the amount of  $\text{H}^+$  is only significant from the first step. I'll explain this better as we go through the next Example.

.....

**Example 6.** 0.00203 mol  $\text{CO}_2$  is dissolved in water to make 1.00 L of solution. What are the concentrations of  $\text{CO}_2$ ,  $\text{H}^+$  and  $\text{HCO}_3^-$  at equilibrium? What is the pH of the solution?

.....

Consider again the two dissociation steps.



Note that the problem asks for concentrations of  $\text{CO}_2$ ,  $\text{H}^+$  and  $\text{HCO}_3^-$ . All of these are directly involved in the first step.  $\text{H}^+$  and  $\text{HCO}_3^-$  are also involved in the second step, but we will wait on that momentarily and do the first step first.

$$K_{a1} = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{CO}_2]} = 4.5 \times 10^{-7}$$

Initial  $[\text{CO}_2]$  is 0.00203 M. Now, set up a table.

	$[\text{CO}_2]$	$[\text{H}^+]$	$[\text{HCO}_3^-]$
Initial:	0.00203	-0-	-0-
Changes:	-x	+x	+x
Equilibrium:	0.00203 - x	x	x

Plug in.

$$K_{a1} = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{CO}_2]} = \frac{(x)(x)}{0.00203 - x} = 4.5 \times 10^{-7}$$

How do you wish to solve?  $K_{a1}$  is very small which is good for approximation, but the sample is very dilute. Approximation/iteration? Quadratic? All of the above? Sure, why not?

First, assume  $0.00203 - x \approx 0.00203$ .

$$K_{a1} = \frac{(x)(x)}{0.00203 - x} \approx \frac{x^2}{0.00203} = 4.5 \times 10^{-7}$$

This gives  $x = 3.0 \times 10^{-5}$ . So far, that is simple approximation. How good an answer is this? Iterate.

$$K_{a1} \approx \frac{(x)(x)}{0.00203 - 0.000030} = \frac{x^2}{0.00200} = 4.5 \times 10^{-7}$$

You will again get  $x = 3.0 \times 10^{-5}$ . Stop.

For the quadratic method, start from  $K_{a1}$

$$K_{a1} = \frac{(x)(x)}{0.00203 - x} = 4.5 \times 10^{-7}$$

and re-arrange to

$$x^2 + 4.5 \times 10^{-7} x - 9.1 \times 10^{-10} = 0$$

and plug into quadratic to get  $x = 3.0 \times 10^{-5}$ , same as above.

This gives our answers for first dissociation.

$$[\text{CO}_2] = (0.00203 - 3.0 \times 10^{-5}) \text{ M} = 0.00200 \text{ M}$$

$$[\text{H}^+] = [\text{HCO}_3^-] = 3.0 \times 10^{-5} \text{ M}$$


$$\text{pH} = -\log(3.0 \times 10^{-5}) = 4.52$$

That ends this Example as originally stated. But what about the second dissociation?

Although additional  $\text{H}^+$  is produced in the second dissociation, the self-imposed simplification says that the additional amount is not significant compared to the amount from the first step. Ergo, we're done. The second step will also affect the amount of  $\text{HCO}_3^-$ , but that effect will likewise not be significant to the first step. Thus, we work only with the amounts of  $\text{H}^+$  and  $\text{HCO}_3^-$  from the first step. This approach is valid whenever the value of  $K_{a2}$  is very small compared to the value of  $[\text{H}^+]$  from the first step, which is certainly the case in this Example.

OK, so we've got  $[\text{CO}_2]$ ,  $[\text{H}^+]$  and  $[\text{HCO}_3^-]$  as requested, but what if we also needed to find  $[\text{CO}_3^{2-}]$ ?  $\text{CO}_3^{2-}$  is not in the first step. If you need to find the concentration for carbonate, then you have no choice but to invoke the second step. We will now extend this Example to ask this very question: for the solution as given, what is the concentration of  $\text{CO}_3^{2-}$  at equilibrium?

Bring in the second step.



$$K_{a2} = \frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]} = 4.7 \times 10^{-11}$$

You will solve for  $[\text{CO}_3^{2-}]$  in the usual way. First, we need initial  $[\text{HCO}_3^-]$ . The initial amounts for the second step pick up where the first step left off. Thus, initial  $[\text{HCO}_3^-] = 3.0 \times 10^{-5} \text{ M}$ . We also have some initial  $[\text{H}^+]$  going into the second step. Set up a table and spell everything out. I'll use  $y$ 's here to avoid confusion with  $x$ 's in the prior table.

	$[\text{HCO}_3^-]$	$[\text{H}^+]$	$[\text{CO}_3^{2-}]$
Initial:	$3.0 \times 10^{-5}$	$3.0 \times 10^{-5}$	0-
Changes:	$-y$	$+y$	$+y$
Equilibrium:	$3.0 \times 10^{-5} - y$	$3.0 \times 10^{-5} + y$	$y$


Plug in.

$$K_{a2} = \frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]} = \frac{(3.0 \times 10^{-5} + y)(y)}{3.0 \times 10^{-5} - y} = 4.7 \times 10^{-11}$$

This equation is more complicated than usual. Approximate? The  $\text{HCO}_3^-$  is very dilute at  $3.0 \times 10^{-5}$ , but  $K_{a2}$  is  $10^{-11}$ . With such an extremely small  $K_a$ , assume  $y$  is small relative to  $3.0 \times 10^{-5}$ ; this assumption applies for both the numerator and the denominator.

$$K_{a2} = \frac{(3.0 \times 10^{-5} + y)(y)}{3.0 \times 10^{-5} - y} \approx \frac{(3.0 \times 10^{-5})(y)}{3.0 \times 10^{-5}} = 4.7 \times 10^{-11}$$

This gives  $y = 4.7 \times 10^{-11}$ . So far, that's simple approximation. What do you think? Is that a good enough answer? What would happen if you iterate? Well, if you plug that value into the numerator and denominator, how will  $y = 4.7 \times 10^{-11}$  compare to  $3.0 \times 10^{-5}$ ? The value of  $y$  is way too small to make nary a dent in  $3.0 \times 10^{-5}$ ; thus, the values in the numerator and denominator would not change with  $y$ . If you iterate, you will get the same number and that means you're done. The second dissociation step ends as follows.



The concentrations of  $\text{H}^+$  and  $\text{HCO}_3^-$  remain  $3.0 \times 10^{-5} \text{ M}$  as determined for the first step; they are unchanged in the second step to the allowed sigfigs. This will be true within our limit of coverage. A consequence of this coverage is that  $y = K_{a2}$  as is seen in this last calculation. If you remember this, you won't even need to set up for a calculation for the second dissociation.


This ends the polyprotic example with all steps weak. Now let's go strong.

.....

**Example 7.** 0.0200 mol  $\text{H}_2\text{SO}_4$  is dissolved in 1.00 L solution. What are  $[\text{H}^+]$ ,  $[\text{HSO}_4^-]$ ,  $[\text{SO}_4^{2-}]$  and pH at equilibrium?

.....

For  $\text{H}_2\text{SO}_4$ , the first step is strong but the second step is weak. For these cases, you must work with both steps. We begin with the first.



This is strong, so everything goes to the right as is done for any strong acid. The initial 0.0200 M  $\text{H}_2\text{SO}_4$  goes completely to 0.0200 M  $\text{H}^+$  and 0.0200 M  $\text{HSO}_4^-$ . This much is automatic for a strong step.

For the second step, we do the usual equilibrium calculation.



$$K_{a2} = \frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} = 0.010$$

Set up a table. As done in the follow-up to Example 6, the initial amounts for the second step come from the final amounts for the first step.

	$[\text{HSO}_4^-]$	$[\text{H}^+]$	$[\text{SO}_4^{2-}]$
Initial:	0.0200	0.0200	0
Changes:	-x	+x	+x
Equilibrium:	$0.0200 - x$	$0.0200 + x$	x


Plug in.

$$K_{a2} = \frac{[\text{H}^+][\text{SO}_4^{2-}]}{[\text{HSO}_4^-]} = \frac{(0.0200 + x)(x)}{0.0200 - x} = 0.010$$

Let's go through quadratic on this one. Re-arrange

$$\begin{aligned} 0.0200x + x^2 &= 0.000200 - 0.010x \\ x^2 + 0.030x - 0.000200 &= 0 \end{aligned}$$

and then plug into the quadratic equation to get  $x = 0.0056$ . This  $x$  then goes into the bottom line of the table to give the final concentrations.



$K$  check gives 0.010, right on the money. Finally,

$$\text{pH} = 1.59$$

and it's over.

Could we have approximated this one? With a  $K_{a2}$  of 0.010, this is very risky. But, yes, you can derive the same value of  $x$  and it will only take ten iteration steps beyond the initial approximation. Try it if you'd like, but watch your sigfigs.

These Examples 6 and 7 are fairly typical of a dissociation problem for a polyprotic acid. Polyprotic problems can look intimidating, but each type involves only one weak equilibrium calculation. Here it is in summary.

For a polyprotic with all weak steps, you only do a full equilibrium calculation for the first dissociation. The next dissociation step is ignored unless you need its product anion, in which case its concentration has the same value as  $K_{a2}$ .

For a polyprotic whose first step is strong, you simply send the first step all the way to full dissociation, and you do a weak calculation starting from there for the second step.


Our Examples were limited to diprotics. You can do triprotics along a similar approach; all common triprotic acids have all weak steps, so you would follow the method of Example 6.

This concludes our acid dissociation calculations. Before moving on, let's consider a different kind of issue. The acids cited thus far have covered a range of  $K_a$  values, from strong all the way down to  $K_a = 1.0 \times 10^{-10}$  for phenol, and even lower for hydrogen anions such as  $\text{HPO}_4^{2-}$ . We will now take a brief look at why there is such a range.

#### 55.4 Strong, weak and weaker

The thermodynamic parameters of weak acid dissociation were mentioned for acetic acid in Section 54.1, for which  $\Delta H^\circ = -0.25$  kJ and  $\Delta S^\circ = -92.1$  J/K. In general, enthalpies for acid dissociation can run negative or positive. Acids with  $K_a$  values in the range of  $10^{-3}$  to  $10^{-5}$  can go either way, exothermic or endothermic. Weaker acids ( $K_a < 10^{-5}$ ) typically have endothermic dissociation, while acids not as weak ( $K_a > 10^{-3}$ ) tend to run exothermic. Thus,  $\Delta H^\circ$  can favor or oppose dissociation.  $\Delta S^\circ$ , however, will be negative and oppose dissociation for all acids, as was also noted in Section 54.1.

Let's look a bit more at the enthalpy contribution,  $\Delta H^\circ$ , for dissociation in general.



As we have seen in the past, enthalpies of reactions commonly derive from bond energies and from IFs. Let's consider some of the factors which can be involved in acid dissociation. An H-A bond is broken, and there will always be some cost associated with that. An O-H bond is formed within protonated water ( $\text{H}^+(aq)$ ), and that releases some energy. (This  $\text{H}^+$  part is the same for all acids.) There's a large change in hydration energies: hydration of  $\text{A}^-$  will be greater than hydration for HA due to ion-dipole interactions, and that will release some energy. Hydrogen bonding can also contribute, and this may differ for  $\text{HA}(aq)$  vs.  $\text{A}^-(aq)$ . How these, and potentially other, factors compare will determine the sign and the magnitude of the  $\Delta H^\circ$  for the overall dissociation.

As always, it is the specific combination of  $\Delta H^\circ$  and  $\Delta S^\circ$  which determines a specific  $\Delta G^\circ$ , which then determines a specific  $K$ . Notably for acid dissociations, the values of  $\Delta G^\circ$  don't really change by large amounts, but that still has a sizeable effect on  $K_a$ . This goes back to a point made in Section 51.2.

“ The value of  $K$  is very sensitive to the value of  $\Delta G^\circ$  due to their exponential relationship.

$$K = e^{-\Delta G^\circ/RT} \text{ ”}$$

Back then, a table was given for  $\Delta G^\circ$ 's varying by hundreds of kJ's in  $\pm$  directions, and the respective  $K$ 's varied by huge numbers. Now in the case of acid dissociation, values of  $\Delta G^\circ$  will span only tens of kJ's, but even this will affect  $K_a$  substantially. Consider a new table covering some dissociation values for  $\Delta G^\circ$ .

$\Delta G^\circ$ (kJ)	34.2	28.5	22.8	17.1	11.4	...	-11.4	-17.1
$K_a$	$10^{-6}$	$10^{-5}$	$10^{-4}$	$10^{-3}$	$10^{-2}$		$10^2$	$10^3$

Here, each column of  $\Delta G^\circ$  changes by a mere 5.7 kJ, which is small. Nevertheless, each 5.7 kJ translates into a factor of 10 for  $K_a$ , which is sizeable. Thus, small changes in chemical properties between different conjugate pairs can have a large effect on  $K_a$ . Note the cases of negative  $\Delta G^\circ$  in the table; these apply to strong acids. Although we tend to equate all strong acids as equally strong in water, they still differ in  $\Delta G^\circ$ .

So what kind of chemical properties have an impact on acidity? That depends.

There are many different categories of acids, and comparisons across the categories are not straightforward. Within a category, however, trends can sometimes be identified which indicate why some acids are better or weaker than others, but even these are subject to subtleties of  $\Delta H^\circ$  and  $\Delta S^\circ$ . We will consider two trends.

The first trend involves polyprotic acids: each successive dissociation is harder to do. Thus,  $K_{a2} < K_{a1}$  and  $K_{a3} < K_{a2}$  for a particular acid. The reason for this is as follows. Consider  $\text{H}_3\text{PO}_4$  again.



After each step, you end with a more negative ion. It's harder to pull  $\text{H}^+$  off a more negative ion, so the next step costs more in enthalpy. Also for each step, the more negative ion is worse for entropy. Overall, each step is more and more opposed by  $\Delta H^\circ$  and  $\Delta S^\circ$ , and each step is weaker.

The second trend is for oxyacids. Oxyacids contain a central atom bonded to one or more oxygens, one or more of which are bonded to hydrogen. A generic formula is  $(\text{HO})_x\text{ZO}_y$  with  $x \geq 1$  and  $y \geq 0$ . There are many such acids and many of these are very common; among these are  $\text{H}_2\text{SO}_4$ ,  $\text{H}_3\text{PO}_4$  and  $\text{HNO}_3$ , which are the top three acids produced industrially worldwide. Other examples include  $\text{H}_2\text{CO}_3$ ,  $\text{HClO}_4$ ,  $\text{HClO}_3$ , etc. In all of these here, the H's are bonded to O's, despite the way the formula is written. For example,  $\text{H}_2\text{SO}_4$  can be written  $(\text{HO})_2\text{SO}_2$  while  $\text{HClO}_3$  can be written  $\text{HOClO}_2$ . For oxyacids in general, there are two comparisons to note for acid strength. First, for one specific element Z, the  $K_a$  increases (better acid) when more O's are present. More O's present also means the oxidation number of Z is higher, so we can also say that  $K_a$  increases as the oxidation number of Z increases. As an illustration, there are four oxyacids for Cl and their strengths run  $\text{HClO}_4 > \text{HClO}_3 > \text{HClO}_2 > \text{HClO}$ ; the ON's for Cl in that series run 7, 5, 3, 1. Second, for a specific formula but different element Z, then  $K_a$  increases as the electronegativity of Z increases. For example, the strengths for the "hypohalous" acid series,  $\text{HZO}$  with Z = halogen, runs  $\text{HClO} > \text{HBrO} > \text{HIO}$ .

With this, we pause on acids temporarily while we bring in some basic material.

## Problems

- True or false.
  - In a weak acid solution, dilution favors dissociation.
  - Perchloric acid is diprotic.
  - Water can deprotonate bisulfate ion (at least to some extent).
  - The  $K_a$  of  $\text{H}_2\text{PO}_4^-$  is greater than the  $K_a$  of  $\text{HPO}_4^{2-}$ .
  - $\text{HClO}_2$  is a stronger acid than  $\text{HClO}_3$ .
- Write the balanced equation for acid dissociation and write the  $K_a$  expression for each of the following.
  - $\text{HBrO}$
  - $\text{HN}_3$
- Write the balanced equations for each step of dissociation for arsenic acid,  $\text{H}_3\text{AsO}_4$ .
- Which of the following acids are monoprotic?
 

nitric acid      phosphoric acid      oxalic acid      chlorous acid      carbonic acid
- Pyrophosphoric acid,  $\text{H}_4\text{P}_2\text{O}_7$ , is tetraprotic, meaning it is capable of losing four  $\text{H}^+$ . Write the balanced equation and the  $K_a$  expression for the fourth step.
- A 150. mL solution is prepared using 0.0198 mol butyric acid,  $\text{C}_3\text{H}_7\text{CO}_2\text{H}$ . What are the concentrations of  $\text{C}_3\text{H}_7\text{CO}_2\text{H}$ ,  $\text{H}^+$  and  $\text{C}_3\text{H}_7\text{CO}_2^-$  at equilibrium? What are the pH and the percent dissociation?

- 0.0268 mol of chloroacetic acid,  $\text{ClCH}_2\text{CO}_2\text{H}$ , is dissolved in water to make 800. mL of solution. What are the concentrations of  $\text{ClCH}_2\text{CO}_2\text{H}$ ,  $\text{H}^+$  and  $\text{ClCH}_2\text{CO}_2^-$ ? What are the pH and the percent dissociation at equilibrium?
- Vinegar is a mixture of various components, and the dominant acid component (and also the dominant smell) is acetic acid. The acid content can vary, typically around 5% w/w. As a comparable sample, consider a solution which contains 5.14 g  $\text{CH}_3\text{CO}_2\text{H}$  dissolved in water to give 100. mL solution. What is the pH of this solution?
- Vitamin C is ascorbic acid,  $\text{C}_6\text{H}_8\text{O}_6$ , which is a diprotic acid, weak in both steps. 0.0112 mol of ascorbic acid is contained within 500. mL of solution. The pH of the solution is 2.92 at equilibrium. Find  $K_a$  and  $\Delta G^\circ$  for the first dissociation step at 25 °C.
- A solution is prepared using 0.124 mol  $\text{H}_2\text{C}_2\text{O}_4$  in a volume of 1.00 L. At equilibrium, what are the concentrations of  $\text{H}_2\text{C}_2\text{O}_4$ ,  $\text{H}^+$  and  $\text{HC}_2\text{O}_4^-$ ? What is the percent dissociation of  $\text{H}_2\text{C}_2\text{O}_4$ ? What is the concentration of  $\text{C}_2\text{O}_4^{2-}$ ?
- Like sulfuric acid, selenic acid is also strong in the first step and weak in the second;  $K_{a2} = 0.018$ . A 200. mL solution of  $\text{H}_2\text{SeO}_4$  is prepared using 0.00740 mol  $\text{H}_2\text{SeO}_4$ . What is the pH of the solution at equilibrium?
- Of the following acids, which one is the strongest? Which one is the weakest?  
 $\text{HClO}$        $\text{HClO}_2$        $\text{HClO}_3$        $\text{HBrO}_3$        $\text{HBrO}_2$        $\text{HIO}$
- Consider the acid dissociation of hydrocyanic acid. Starting from values for  $\Delta H_f^\circ$  and for  $S^\circ$ , calculate  $\Delta H^\circ$  and  $\Delta S^\circ$  for the dissociation. Is dissociation endothermic or exothermic? Will the  $K_a$  be greater at 25 °C or at 70. °C? Calculate  $\Delta G^\circ$  at 70. °C. Use that to find  $K_a$  at 70. °C and compare that to the value at 25 °C.
- Calculate  $[\text{H}^+]$  and the pH of carbonated water when the pressure of  $\text{CO}_2(g)$  above the solution at 25 °C at equilibrium is 4.0 atm. (Here's a start. Set up the solubility equilibrium for  $\text{CO}_2(g)$  and also set up the first dissociation step for  $\text{CO}_2(aq)$ . Add these two equations together. Find  $\Delta G^\circ$  for the sum equation from free energies of formation and use that to find  $K$ .)